

Occult Scaphoid Fractures: A Multi-Centre Service Evaluation of MRI Practice

WhichAudit · NHS Clinical Audit Repository · [whicaudit.com](https://www.whicaudit.com)

Type	AUDIT
Specialty	orthopaedics
Trust	Guy's and St Thomas' NHS Foundation Trust
Year	2022
Added	25 Apr 2026

Background

Occult scaphoid fractures — those not visible on initial plain radiographs but subsequently detected on MRI — represent a clinically important diagnostic challenge. Missed or delayed diagnosis can result in avascular necrosis and non-union, with significant long-term morbidity. This multi-centre evaluation assessed the rate of complications and subsequent surgical intervention in patients with occult scaphoid fractures identified via MRI at GSTT and collaborating centres.

Aims

To assess the complications arising from occult scaphoid fractures detected by early MRI at GSTT, and to evaluate the outcomes of this diagnostic approach across collaborating centres.

Objectives

1. Retrospectively identify all patients who underwent wrist MRI following a normal scaphoid series between January and December 2022. 2. Document the proportion in whom an occult scaphoid fracture was confirmed on MRI. 3. Record rates of avascular necrosis, non-union and requirement for operative intervention. 4. Compare complication rates against published literature benchmarks.

Guidelines

NICE CG 102 / British Society for Surgery of the Hand guidelines on scaphoid fractures

Keywords

scaphoid fracture, occult fracture, MRI, wrist, avascular necrosis, hand surgery